

# VACCINE INFORMATION STATEMENT

## Influenza (Flu) Vaccine (Inactivated or Recombinant)

### What You Need to Know

Many Vaccine Information Statements are available in Spanish and other languages. See [www.immunize.org/vis](http://www.immunize.org/vis)

Hojas de información sobre vacunas están disponibles en español y en muchos otros idiomas. Visite [www.immunize.org/vis](http://www.immunize.org/vis)

### 1. Why get vaccinated?

**Influenza vaccine can prevent influenza (flu).**

**Flu** is a contagious disease that spreads around the United States every year, usually between October and May. Anyone can get the flu, but it is more dangerous for some people. Infants and young children, people 65 years and older, pregnant people, and people with certain health conditions or a weakened immune system are at greatest risk of flu complications.

Pneumonia, bronchitis, sinus infections and ear infections are examples of flu-related complications. If you have a medical condition, such as heart disease, cancer or diabetes, flu can make it worse.

Flu can cause fever and chills, sore throat, muscle aches, fatigue, cough, headache, and runny or stuffy nose. Some people may have vomiting and diarrhea, though this is more common in children than adults.

Each year **thousands of people in the United States die from flu**, and many more are hospitalized. Flu vaccine prevents millions of illnesses and flu-related visits to the doctor each year.

### 2. Influenza vaccines

CDC recommends everyone 6 months and older get vaccinated every flu season. **Children 6 months through 8 years of age** may need 2 doses during a single flu season. **Everyone else** needs only 1 dose each flu season.

It takes about 2 weeks for protection to develop after vaccination.

There are many flu viruses, and they are always changing. Each year a new flu vaccine is made to protect against the influenza viruses believed to be likely to cause disease in the upcoming flu season. Even when the vaccine doesn't exactly match these viruses, it may still provide some protection.

Influenza vaccine **does not cause flu**.

Influenza vaccine may be given at the same time as other vaccines.

### 3. Talk with your health care provider

Tell your vaccine provider if the person getting the vaccine:

- Has had an **allergic reaction after a previous dose of influenza vaccine**, or has any **severe, life-threatening allergies**.
- Has ever had **Guillain-Barré Syndrome** (also called GBS).

In some cases, your health care provider may decide to postpone influenza vaccination until a future visit.

Influenza vaccine can be administered at any time during pregnancy. People who are or will be pregnant during influenza season should receive inactivated influenza vaccine.

People with minor illnesses, such as a cold, may be vaccinated. People who are moderately or severely ill should usually wait until they recover before getting influenza vaccine.

Your health care provider can give you more information.

### 4. Risks of a vaccine reaction

- Soreness, redness, and swelling where shot is given, fever, muscle aches, and headache can happen after influenza vaccination.
- There may be a very small increased risk of Guillain-Barré Syndrome (GBS) after inactivated influenza vaccine (the flu shot).

Young children who get the flu shot along with pneumococcal vaccine (PCV13), and/or DTaP vaccine at the same time might be slightly more likely to have a seizure caused by fever. Tell your health care provider if a child who is getting flu vaccine has ever had a seizure.

People sometimes faint after medical procedures, including vaccination. Tell your provider if you feel dizzy or have vision changes or ringing in the ears.

As with any medicine, there is a very remote chance of a vaccine causing a severe allergic reaction, other serious injury, or death.

### 5. What if there is a serious problem?

An allergic reaction could occur after the vaccinated person leaves the clinic. If you see signs of a severe allergic reaction (hives, swelling of the face and throat, difficulty breathing, a fast heartbeat, dizziness, or weakness), call **9-1-1** and get the person to the nearest hospital.

For other signs that concern you, call your health care provider.

Adverse reactions should be reported to the Vaccine Adverse Event Reporting System (VAERS). Your health care provider will usually file this report, or you can do it yourself. Visit the VAERS website at [www.vaers.hhs.gov](http://www.vaers.hhs.gov) or call **1-800-822-7967**. *VAERS is only for reporting reactions, and VAERS staff do not give medical advice.*

### 6. The National Vaccine Injury Compensation Program

The National Vaccine Injury Compensation Program (VICP) is a federal program that was created to compensate people who may have been injured by certain vaccines. Claims regarding alleged injury or death due to vaccination have a time limit for filing, which may be as short as two years. Visit the VICP website at [www.hrsa.gov/vaccinecompensation](http://www.hrsa.gov/vaccinecompensation) or call **1-800-338-2382** to learn about the program and about filing a claim.

### 7. How can I learn more?

- Ask your healthcare provider.
- Call your local or state health department
- Visit the website of the Food and Drug Administration (FDA) for vaccine package inserts and additional information at [www.fda.gov/vaccines-blood-biologics/vaccines](http://www.fda.gov/vaccines-blood-biologics/vaccines)
- Contact the Centers for Disease Control and Prevention (CDC)
  - Call **1-800-232-4636 (1-800-CDC-INFO)**
  - Visit CDC's website at [www.cdc.gov/flu](http://www.cdc.gov/flu)

42 U.S.C. § 300aa-26

8/6/2021

OFFICE  
USE  
ONLY



U.S. Department of  
Health and Human Services  
Centers for Disease  
Control and Prevention

Healthy Solutions, Inc. 913-345-2223

# VACCINE INFORMATION STATEMENT

## Tdap Vaccine (Tetanus, Diphtheria, and Pertussis) What you need to know

Many Vaccine Information Statements are available in Spanish and other languages. See [www.immunize.org/vis](http://www.immunize.org/vis)

Hojas de información sobre vacunas están disponibles en español y en muchos otros idiomas. Visite [www.immunize.org/vis](http://www.immunize.org/vis)

### 1. Why get vaccinated?

Tdap vaccine can prevent **tetanus, diphtheria, and pertussis**.

Diphtheria and pertussis spread from person to person. Tetanus enters the body through cuts or wounds.

- **TETANUS (T)** causes painful stiffening of the muscles. Tetanus can lead to serious health problems, including being unable to open the mouth, having trouble swallowing and breathing, or death.
- **DIPHTHERIA (D)** can lead to difficulty breathing, heart failure, paralysis, or death.
- **PERTUSSIS (aP)**, also known as “whooping cough,” can cause uncontrollable, violent coughing which makes it hard to breathe, eat, or drink. Pertussis can be extremely serious especially in babies and young children, causing pneumonia, convulsions, brain damage, or death. In teens and adults, it can cause weight loss, loss of bladder control, passing out, and rib fractures from severe coughing.

### 2. Tdap vaccine

Tdap is only for children 7 years and older, adolescents, and adults.

**Adolescents** should receive a single dose of Tdap, preferably at age 11 or 12 years.

**Pregnant people** should get a dose of Tdap during every pregnancy, preferably during the early part of the third trimester, to help protect the newborn from pertussis. Infants are most at risk for severe, life-threatening complications from pertussis.

**Adults** who have never received Tdap should get a dose of Tdap.

Also, **adults should receive a booster dose of either Tdap or Td** (a different vaccine that protects against tetanus and diphtheria but not pertussis) **every 10 years**, or after 5 years in the case of a severe or dirty wound or burn.

Tdap may be given at the same time as other vaccines.

### 3. Talk with your health care provider

Tell your vaccine provider if the person getting the vaccine:

- Has had an **allergic reaction after a previous dose of any vaccine that protects against tetanus, diphtheria, or pertussis**, or has any severe, life-threatening allergies.
- Has had a **coma, decreased level of consciousness, or prolonged seizures within 7 days after a previous dose of any pertussis vaccine (DTP, DTaP, or Tdap)**.
- Has **seizures or another nervous system problem**.
- Has ever had **Guillain-Barré Syndrome** (also called GBS).
- Has had **severe pain or swelling after a previous dose of any vaccine that protects against tetanus or diphtheria**.

In some cases, your health care provider may decide to postpone Tdap vaccination to a future visit.

People with minor illnesses, such as a cold, may be vaccinated. People who are moderately or severely ill should usually wait until they recover before getting Tdap vaccine.

Your health care provider can give you more information.

### 4. Risks of vaccine reaction

- Pain, redness, or swelling where the shot was given, mild fever, headache, feeling tired, and nausea, vomiting, diarrhea, or stomachache sometimes happen after Tdap vaccination.

People sometimes faint after medical procedures, including vaccination. Tell your provider if you feel dizzy or have vision changes or ringing in the ears.

As with any medicine, there is a very remote chance of a vaccine causing a severe allergic reaction, other serious injury, or death.

### 5. What if there is a serious problem?

An allergic reaction could occur after the vaccinated person leaves the clinic. If you see signs of a severe allergic reaction (hives, swelling of the face and throat, difficulty breathing, a fast heartbeat, dizziness, or weakness), call **9-1-1** and get the person to the nearest hospital.

For other signs that concern you, call your health care provider.

Adverse reactions should be reported to the Vaccine Adverse Event Reporting System (VAERS). Your health care provider will usually file this report, or you can do it yourself. Visit the VAERS website at [www.vaers.hhs.gov](http://www.vaers.hhs.gov) or call **1-800-822-7967**. *VAERS is only for reporting reactions, and VAERS staff do not give medical advice.*

### 6. The National Vaccine Injury Compensation Program

The National Vaccine Injury Compensation Program (VICP) is a federal program that was created to compensate people who may have been injured by certain vaccines. Claims regarding alleged injury or death due to vaccination have a time limit for filing, which may be as short as two years. Visit the VICP website at [www.hrsa.gov/vaccinecompensation](http://www.hrsa.gov/vaccinecompensation) or call **1-800-338-2382** to learn about the program and about filing a claim.

### 7. How can I learn more?

- Ask your health care provider.
- Call your local or state health department.
- Visit the website for the Food and Drug Administration (FDA) for vaccine package inserts and additional information at [www.fda.gov/vaccines-blood-biologics/vaccines](http://www.fda.gov/vaccines-blood-biologics/vaccines)
- Contact the Centers for Disease Control and Prevention (CDC):
  - Call **1-800-232-4636 (1-800-CDC-INFO)** or
  - Visit CDC's website at [www.cdc.gov/vaccines](http://www.cdc.gov/vaccines)

42 U.S.C. § 300aa-26

8/6/2021

OFFICE  
USE  
ONLY



U.S. Department of  
Health and Human Services  
Centers for Disease  
Control and Prevention

Healthy Solutions, Inc. 913-345-2223